arXiv:0806.0499v1 [hep-ph] 3 Jun 2008

QED peripheral mechanism of pair production at colliders

A. I. Ahmadov*
Joint Institute for Nuclear Research, Dubna, Russia and
Institute of Physics, Azerbaijan National Academy of Sciences, Baku, Azerbaijan

M. V. Galynskii†
Institute of Physics, Minsk, Belorussia

Yu. M. Bystritskiy‡ and E. A. Kuraev§
Joint Institute for Nuclear Research, Dubna, Russia

M. G. Shatnev¶
Kharkov Institute of Phys. and Tech., Kharkov, 61108, Ukraine

Cross section of the processes of neutral pion production as well as pairs of charged fermions and bosons in peripherical interaction of leptons, photons are calculated in main logarithmical approximation. We investigate the phase volumes and differential cross sections. The differential cross section of few neutral pions and a few pairs production are written down explicitly. Considering the academic problem of summation on the number of pairs for the case of massless particles we reproduce the known results obtained in seventies of last century. The possibility to construct the generator for Monte-Carlo modeling of this processes basing of this results are discussed.

I. INTRODUCTION

The motivation of this paper is the need of creation of realistic Monte-Carlo generators describing the multiparticle production at high energy colliders. We consider the case of peripherical interaction of initial particles which corresponds to the kinematics of final particles which group a set of jets flying in the wide range of rapidities – starting from fragmentation region up to pionization region.

Keeping in mind the possibility to construct the $\gamma - e$ and $\gamma - \gamma$ colliders on the basis of the e^+e^--colliders using the backwards scattering of laser beam we consider the processes which may be relevant for experiments on this facilities.

Formulae, obtained below can be applied to the case of γp, ep and $p(\bar{p})p$ colliders with a straightforward generalization.

Processes of charged pairs production at high energy beams $a - b$ collisions where $a, b = \gamma, e^\pm$ have a set of general properties. To remind them let consider process of e^+e^- pair production at high energy $e^\pm e^-$ collisions in peripheral kinematics. Matrix

---

*Electronic address: Azad.Ahmedov@sunse.jinr.ru
†Electronic address: galynski@dragon.bas-net.by
‡Electronic address: bystr@theor.jinr.ru
§Electronic address: kuraev@theor.jinr.ru
¶Electronic address: mshatnev@yahoo.com



FIG. 1: The two-photon mechanism of production of a pion pair.

element of production of a pair of charged particles in high energy electron-positron collisions (center of mass of initial colliding beams is implied)

$$e^+(p) + e^-(k) \to e^+(p') + e^-(k') + q(q_-) + \bar{q}(q_+), \tag{1}$$

$$s = 2pk \gg -q_1^2 = -(k-k')^2 \sim -q_2^2 = (p-p')^2 \sim q_\pm^2 = m^2, \tag{2}$$

$$k^2 = m_1^2, \qquad p^2 = m_2^2, \tag{3}$$

have a form (here and further we consider the so called «two-photon mechanism» of pair production) (see Fig. 1):

$$M^i = \frac{(4\pi\alpha)^2}{q_1^2 q_2^2} \left[\bar{u}(p')\gamma^\mu u(p)\right] \left[\bar{v}(k')\gamma^\nu v(k)\right] O^i_{\mu_1\nu_1} g_{\mu\mu_1} g_{\nu\nu_1}, \qquad i = e, \pi. \tag{4}$$

We use Gribov representation for photon Green function

$$q_{\mu\nu} = q^\perp_{\mu\nu} + \frac{1}{s}\left(p_\mu k_\nu + p_\nu k_\mu\right) \tag{5}$$

and Sudakov decomposition of 4-momenta

$$\begin{aligned} &q_1 = \beta_1 \tilde{k} + q_1^\perp, \qquad q_2 = \alpha_2 \tilde{p} + q_2^\perp, \\ &q_\pm = \alpha_\pm \tilde{p} + \beta_\pm \tilde{k} + q_\pm^\perp, \qquad s\alpha_\pm \beta_\pm = \vec{q}_\pm^{\,2} + m^2. \end{aligned} \tag{6}$$

over «almost light-cone» 4-vectors $\tilde{p} = p - k(m_2^2/s)$, $\tilde{k} = k - p(m_1^2/s)$. Further the tilde subscripts will be omitted.

Keeping in mind the conservation law, $q_1 + q_2 = q_+ + q_-$ and the hierarchy of Sudakov parameters, $\beta_1 \gg \beta_2$, $\alpha_1 \ll \alpha_2$, the matrix element (4) can be written down in the form

$$M^i = s\frac{(8\pi\alpha)^2}{q_1^2 q_2^2} N_1 N_2 \Phi^i(q_1, q_2), \qquad i = e, \pi \tag{7}$$

$$N_1 = \frac{1}{s}\bar{v}(k')\hat{p}v(k), \qquad N_2 = \frac{1}{s}\bar{u}(p')\hat{k}u(p), \tag{8}$$

$$\sum |N_1|^2 = \sum |N_2|^2 = 2, \tag{9}$$

where factor $\Phi^i(q_1, q_2) = \Phi^i_{12} = (1/s)k^\mu p^\nu O^i_{\mu\nu}$ is the light-cone projection of amplitudes of the subprocess of charged particles pair production from two virtual photons: $\gamma^*(q_1)\gamma^*(q_2) \to q(q_-)\bar{q}(q_+)$. It is important to note that the quantity Φ^i_{12} is finite in limit $s \to \infty$ as well as $N_{1,2}$.

II. THE SUBPROCESS $\gamma^*(q_1)\gamma^*(q_2) \to q(q_-)\bar{q}(q_+)$

In case of electron-positron pair production we have:

$$\Phi^e_{12} = \frac{1}{s}\bar{u}(q_-)\left(\hat{k}\frac{\hat{q}_- - \hat{q}_1 + m}{-D_1/x_+}\hat{p} + \hat{p}\frac{-\hat{q}_+ + \hat{q}_1 + m}{-D_2/x_-}\hat{k}\right)v(q_+), \tag{10}$$

where

$$\begin{aligned} D_1 &= -x_+\left((q_- - q_1)^2 - m^2\right) = \vec{q}_+^{\,2} - 2x_+\vec{q}_+\vec{q}_2 + x_+\vec{q}_2^{\,2} + m^2, \\ D_2 &= -x_-\left((q_- - q_2)^2 - m^2\right) = \vec{q}_-^{\,2} - 2x_-\vec{q}_-\vec{q}_2 + x_-\vec{q}_2^{\,2} + m^2, \end{aligned}$$

where $x_\pm$ are the energy fractions of pair components ($x_\pm = \alpha_\pm/\alpha_2$, $x_+ + x_- = 1$) and $\vec{q}_i$ is two-dimensional Euclidean vector. Using Dirac equation $\hat{q}_-u(q_-) = mu(q_-)$, $\hat{q}_+v(q_+) = -mv(q_+)$ we can rewrite formula (10) as:

$$\Phi^e_{12} = \frac{1}{s}\bar{u}(q_-)\left(A\hat{p} + B\hat{k}\hat{q}_1\hat{p} + C\hat{p}\hat{q}_1\hat{k}\right)v(q_+), \tag{11}$$

where

$$A = sx_+x_-\left(\frac{1}{D_2} - \frac{1}{D_1}\right), \qquad B = \frac{x_+}{D_1}, \qquad C = -\frac{x_-}{D_2}.$$

In case of pion pair production performing the same algebra we obtain:

$$\Phi^\pi_{12} = x_+x_-\left(\frac{-2\vec{q}_+\vec{q}_2 + \vec{q}_2^{\,2}}{D_1} + \frac{-2\vec{q}_-\vec{q}_2 + \vec{q}_2^{\,2}}{D_2}\right). \tag{12}$$

It can be checked that both Φ^e_{12}, Φ^π_{12} tends to zero at $\vec{q}_1 \to 0$ or $\vec{q}_2 \to 0$. Below we will use the impact factor for production of charged particles pair with invariant mass $s_1 = (q_+ + q_-)^2$:

$$J^i_{12} = \int \frac{ds_1 d\Gamma_\pm}{\pi}\left|\Phi^i_{12}\right|^2, \qquad i = e, \pi. \tag{13}$$

The phase volume we parameterize as (we use the relation $d^4q_\pm = \frac{s}{2}d\alpha_\pm d\beta_\pm d^2q_{\pm\perp}$)

$$\frac{1}{\pi}ds_1 d\Gamma_\pm = \frac{1}{\pi}ds_1\frac{d^3q_+}{2E_+}\frac{d^3q_-}{2E_-}\delta^4(q_1+q_2-q_+-q_-) = \frac{dx_+d^2q_+}{2\pi x_+x_-}. \tag{14}$$

And performing the phase volume integration we obtain (see details in Appendix C):

$$\begin{aligned} J^e_{12} &= 4\int\limits_0^1 dx\int\limits_0^1\frac{dt}{D}\left[\vec{q}_1^{\,2}\vec{q}_2^{\,2}\left(a+b-4ab\right)-2ab\left(\vec{q}_1\vec{q}_2\right)^2\right], \\ D &= m^2+a\vec{q}_1^{\,2}+b\vec{q}_2^{\,2}, \end{aligned} \tag{15}$$

where $a=t(1-t)$ and $b=x(1-x)$. After averaging over $\vec{q}_1$ and $\vec{q}_2$ vectors orientations we get the well known result [1, 2, 3]:

$$\bar{J}^e_{12} = 4\vec{q}_1^{\,2}\vec{q}_2^{\,2}\int\limits_0^1\frac{dxdt}{D}\left[a+b-5ab\right]. \tag{16}$$

In case of pion pair production we have:

$$\begin{aligned} J^\pi_{12} &= \frac{1}{2}\int\limits_0^1\frac{dxdt}{D}\left[\vec{q}_1^{\,2}\vec{q}_2^{\,2}\left(1-4a\right)\left(1-4b\right)+8ab\left(\vec{q}_1\vec{q}_2\right)^2\right], \\ \bar{J}^\pi_{12} &= \frac{1}{2}\vec{q}_1^{\,2}\vec{q}_2^{\,2}\int\limits_0^1\frac{dxdt}{D}\left[1-4a-4b+20ab\right]. \end{aligned} \tag{17}$$

We also need to consider the sub-process of pair production by one real and one virtual photons $\gamma(k)+\gamma^*(\chi)\to e^+(q_+)+e^-(q_-)$ and $\gamma(k)+\gamma^*(\chi)\to\pi^+(q_+)+\pi^-(q_-)$. The values similar to (10) and (12) in that case reads as:

$$\begin{aligned} \Phi^e_0 &= \frac{1}{s}\bar{u}(q_-)\left(\hat{e}\frac{-\hat{q}_++\hat{\chi}+m}{(q_--k)^2-m^2}\hat{p}+\hat{p}\frac{\hat{q}_--\hat{\chi}+m}{(-q_+-k)^2-m^2}\hat{e}\right)v(p_+), \\ \Phi^\pi_0 &= \frac{1}{s}\left(\frac{(2q_--k)e-(-2q_++\chi)p}{(q_--k)^2-m^2}+\frac{(-2q_++k)e-(2q_--\chi)p}{(-q_++k)^2-m^2}-2ep\right), \end{aligned} \tag{18}$$

where e is the initial photon polarization 4-vector $ee^*=-\vec{e}\vec{e}^{\,*}=-1$, $ep=ek=0$. Again, using Dirac equation we can rewrite them in form:

$$\Phi^e_0 = \bar{u}(q_-)\left[x_+x_-\left(\frac{1}{D_-}-\frac{1}{D_+}\right)\hat{e}-\frac{x_-}{sD_-}\hat{e}\hat{\chi}\hat{p}+\frac{x_+}{sD_+}\hat{p}\hat{\chi}\hat{e}\right]v(q_+), \tag{19}$$

$$\Phi^\pi_0 = -2x_+x_-\left[\frac{\vec{q}_-\vec{e}}{D_-}+\frac{\vec{q}_+\vec{e}}{D_+}\right], \qquad D_\pm=\vec{q}_\pm^{\,2}+m^2. \tag{20}$$

So now we define the values similar to (13) in case of one real photon:

$$I^{eA}_{01} = \int\frac{ds_1d\Gamma_\pm}{\pi}\left|\Phi^e_0\right|^2 = \frac{2}{3}\int\limits_0^1\frac{dt}{D_0}\left(\vec{e}_A\vec{e}^{\,*}_A\vec{\chi}^2(1+2a)-2a(\vec{e}_A\vec{\chi})(\vec{e}^{\,*}_A\vec{\chi})\right), \tag{21}$$

$$I_{01}^{\pi A} = \int \frac{ds_1 d\Gamma_\pm}{\pi} |\Phi_0^\pi|^2 = \frac{1}{6}\int_0^1 \frac{dt}{D_0}\left(\vec{e}_A\vec{e}_A^{\,*}\vec{\chi}^2(1-4a) + 4a(\vec{e}_A\vec{\chi})(\vec{e}_A^{\,*}\vec{\chi})\right), \qquad (22)$$

where $D_0 = m^2 + a\vec{\chi}^2$ and $a = t(1-t)$. The similar expression can be written down for the case when particle B is a photon with polarization vector e_B: $e_A \to e_B$.

Having this boundary impact factors (21) and (22) we can write the cross sections of production of two pairs of charged particles in two polarized photons collision:

$$\sigma_{ij}^{\gamma\gamma} = \frac{\alpha^4}{\pi}\frac{d^2\vec{q}}{(\vec{q}^2)^2} I^j(\vec{q},\vec{e}_2) I^i(\vec{q},\vec{e}_1), \qquad i,j = e,\pi. \qquad (23)$$

This result is in agreement with Cheng-Wu impact picture of high-energy collisions [4, 5, 6]. Different distributions based on (23) was considered in [7, 8].

To obtain this formula we had rearranged the phase volume of final particles introducing new variables – the transferred momentum $q = \alpha p + \beta k + q_\perp$ and invariant masses of pairs $s_1 = s\alpha$, $s_2 = -s\beta$:

$$\begin{aligned} d\Gamma_{2e^+2e^-}^{\gamma\gamma} &= (2\pi)^{-8}\frac{d^3q_{1+}d^3q_{1-}}{2E_{1+}2E_{1-}}\frac{d^3q_{2+}d^3q_{2-}}{2E_{2+}2E_{2-}}\delta^4(k+p-q_{1+}-q_{1-}-q_{2+}-q_{2-}) \times \\ &\times\ d^4q\,\delta^4(p-q-q_{2+}-q_{2-}) = \\ &= \frac{(2\pi)^{-8}}{2s} d\Gamma_{1\pm}ds_1 d\Gamma_{2\pm}ds_2 d^2\vec{q}, \qquad s_1 = s\alpha, \qquad s_2 = -s\beta, \end{aligned} \qquad (24)$$

where we used the known expression $d^4q = (1/(2s))d^2\vec{q}ds_1ds_2$.

In case of unpolarized photons the cross section of production of two electron-positron pairs is

$$\sigma_{2e^+2e^-}^{\gamma\gamma} = \frac{\alpha^4}{\pi m^2} J_e, \qquad (25)$$

where

$$J_e = \frac{4}{9}\int_0^\infty dz \left(\int_0^1 dx \frac{1+x(1-x)}{1+zx(1-x)}\right)^2 = \frac{1}{36}\left[175\zeta_3 - 38\right] \approx 4.78, \qquad (26)$$

where $\zeta_3 = \sum_{k=1}^\infty \frac{1}{k^3} \approx 1.202...$. In case of production of two $\pi^+\pi^-$ pairs we have:

$$\sigma_{2\pi^+2\pi^-}^{\gamma\gamma} = \frac{\alpha^4}{\pi M_\pi^2} J_\pi, \qquad (27)$$

where

$$J_\pi = \frac{1}{36}\int_0^\infty dz \left(\int_0^1 dx \frac{1-2x(1-x)}{1+zx(1-x)}\right)^2 = \frac{1}{144}\left[7\zeta_3 + 10\right] \approx 0.128. \qquad (28)$$

III. NEUTRAL PIONS PRODUCTION

Consider now the neutral pion production in $\gamma\gamma$-collisions. We will obtain below the cross sections of one, two and a few neutral pions production.

For completeness we put here the results for subprocess $\gamma^*\gamma^* \to \pi^0$ and relevant cross sections.

First let us consider the base decay of neutral pion to two real photons $\pi^0 \to 2\gamma$. It's amplitude have a form:

$$M\left(\pi^0 \to \gamma\left(k_1,e_1\right)\gamma\left(k_2,e_2\right)\right) = \frac{\alpha}{\pi f_\pi}\left(k_1e_1k_2e_2\right), \qquad f_\pi = 94 \text{ MeV}, \tag{29}$$

where we used notation $(abcd) = \varepsilon_{\alpha\beta\gamma\delta}a^\alpha b^\beta c^\gamma d^\delta$. This amplitude gives the well known result:

$$\Gamma_{\pi^0\to2\gamma} = \frac{\alpha^2M_\pi^3}{64\pi^3f_\pi^2} = 7.2 \text{ eV}. \tag{30}$$

Thus for process $\gamma^*\gamma^* \to \pi^0$ we obtain:

$$M_{12}\left(\gamma^*\left(q_1,e_1\right)\gamma^*\left(q_2,e_2\right) \to \pi^0\right) = \frac{\alpha}{\pi f_\pi}\left(q_1e_1q_2e_2\right)R_{12}, \tag{31}$$

where we took into account the external momentum dependence as:

$$R_{12} = R_{21} = 2\int\limits_0^1 dx_1 \int\limits_0^{1-x_1} dx_2 \frac{M^2}{M^2 - M_\pi^2x_2x_3 + \vec{q}_1^2x_1x_2 + \vec{q}_2^2x_1x_3}, \tag{32}$$

where $x_3 = 1 - x_1 - x_2$ and M is the constituent quark mass in the fermion triangle loop. The quantity R_{12} is normalized as $R_{12}|_{M^2\gg M_\pi^2,\vec{q}_i^2} = 1$. If one of photons is real, then:

$$M_{01}\left(\gamma^*\left(q_1,e_1\right)\gamma\left(q_2,e_2\right) \to \pi^0\right) = \frac{\alpha}{\pi f_\pi}\left(q_1e_1q_2e_2\right)R_{01},$$
$$R_{01} = R_{10} = 2\int\limits_0^1 dx_1 \int\limits_0^{1-x_1} dx_2 \frac{M^2}{M^2 - M_\pi^2x_2x_3 + \vec{q}_1^2x_1x_2}.$$

So now we can consider the processes

$$\begin{aligned}
\gamma(k,e_1) + \gamma(p,e_2) &\to \pi^0(p_1) + \pi^0(p_2),\\
\gamma(k,e_1) + \gamma(p,e_2) &\to \pi^0(p_1) + \pi^0(p_2) + \pi^0(p_3),\\
\gamma(k,e_1) + \gamma(p,e_2) &\to \pi^0(p_1) + \pi^0(p_2) + \cdots + \pi^0(p_{n+2}).
\end{aligned}$$

The phase volumes of final states are:

$$d\Gamma^{2\pi^0} = \frac{d^3p_1}{2E_1}\frac{d^3p_2}{2E_2}\left(2\pi\right)^{4-3\cdot2}\delta^4\left(p+k-p_1-p_2\right) =$$

$$= (2\pi)^{-2}\,\frac{d^2q}{2s}, \tag{33}$$

$$d\Gamma^{3\pi^0} = \frac{d^3p_1}{2E_1}\frac{d^3p_2}{2E_2}\frac{d^3p_3}{2E_3}\,(2\pi)^{4-3\cdot 3}\,\delta^4\left(p+k-p_1-p_2-p_3\right) =$$

$$= (2\pi)^{-5}\,\frac{d\beta_1}{\beta_1}\frac{d^2q_1d^2q_2}{4s}, \qquad \frac{M_\pi^2}{s} \ll \beta_1 \ll 1, \tag{34}$$

$$d\Gamma^{(n+2)\pi^0} = \frac{d^3p_1}{2E_1}\frac{d^3p_2}{2E_2}\cdots\frac{d^3p_{n+2}}{2E_{n+2}}\,(2\pi)^{4-3\cdot(n+2)}\,\delta^4\left(p+k-p_1-p_2-\cdots-p_{n+2}\right) =$$

$$= (2\pi)^{4-3(n+2)}\left(\prod_{i=1}^{n}\frac{d\beta_i}{\beta_i}\right)\prod_{i=1}^{n+1}d^2q_i\frac{1}{2^{n+1}s},$$

$$\frac{M_\pi^2}{s} \ll \beta_n \ll \beta_{n-1} \ll \cdots \ll \beta_1 \ll 1. \tag{35}$$

The amplitudes of these processes have a form:

$$M^{2\pi^0} = \left(\frac{\alpha}{\pi f_\pi}\right)^2\frac{s}{2}\,[\vec{e}_1\vec{n}]\,[\vec{e}_2\vec{n}]\,R_{0q}^2, \tag{36}$$

$$M^{3\pi^0} = \left(\frac{\alpha}{\pi f_\pi}\right)^3\frac{s}{2}\,[\vec{e}_1\vec{n}_1]\,[\vec{n}_1\vec{n}_2]\,[\vec{n}_2\vec{e}_2]\,R_{01}R_{12}R_{20}, \tag{37}$$

$$M^{(n+2)\pi^0} = \left(\frac{\alpha}{\pi f_\pi}\right)^{n+2}\frac{s}{2}\,[\vec{e}_1\vec{n}_1]\,[\vec{n}_1\vec{n}_2]\cdots[\vec{n}_n\vec{n}_{n+1}]\,[\vec{n}_{n+1}\vec{e}_2]\times$$

$$\times\; R_{01}R_{12}\cdots R_{n,n+1}R_{n+1,0}, \tag{38}$$

$$\vec{n} = \frac{\vec{q}}{|\vec{q}|}, \qquad \vec{n}_i = \frac{\vec{q}_i}{|\vec{q}_i|},$$

$$[\vec{e}\vec{n}]^2 = 1-(\vec{e}\vec{n})\,(\vec{e}^*\vec{n})\,, \qquad [\vec{n}_1\vec{n}_2]^2 = \sin^2\phi_{12}, \tag{39}$$

where ϕ_{12} is the azimuthal angle between 2-vector $\vec{q}_1$ and $\vec{q}_2$. The cross sections then reads as:

$$d\sigma^{\gamma\gamma\to 2\pi^0} = \frac{\alpha^4}{2^8\pi^5 f_\pi^4}\,[\vec{e}_1\vec{n}]^2\,[\vec{e}_2\vec{n}]^2\,R_{0q}^4\frac{d^2q}{\pi}, \tag{40}$$

$$d\sigma^{\gamma\gamma\to 3\pi^0} = \frac{\alpha^6}{2^{12}\pi^9 f_\pi^6}\,[\vec{e}_1\vec{n}_1]^2\,[\vec{n}_1\vec{n}_2]^2\,[\vec{n}_2\vec{e}_2]^2\,R_{01}^2R_{12}^2R_{20}^2L_\pi\frac{d^2q_1d^2q_2}{\pi^2}, \tag{41}$$

$$d\sigma^{\gamma\gamma\to(n+2)\pi^0} = 2^{-4n-8}\pi^{-4n-5}\left(\frac{\alpha^2}{f_\pi^2}\right)^{n+2}[\vec{e}_1\vec{n}_1]^2\,[\vec{n}_1\vec{n}_2]^2\cdots[\vec{n}_n\vec{n}_{n+1}]^2\,[\vec{n}_{n+1}\vec{e}_2]^2\times$$

$$\times\; R_{01}^2R_{12}^2\cdots R_{n,n+1}^2R_{n+1,0}^2\frac{L_\pi^n}{n!}\left(\prod_{i=1}^{n+1}\frac{d^2q_i}{\pi}\right), \tag{42}$$

here $L_\pi = \ln \frac{s}{M_\pi^2}$. We also can consider the processes $\gamma e \to (n\pi^0)e$ and $ee \to ee(n\pi^0)$ in a same manner.

In case of photo-production of $(n+2)$ pions on electron:

$$\gamma(k,e) + e(p) \;\to\; \pi^0(p_1) + \pi^0(p_2) + \cdots + \pi^0(p_{n+2}) + e(p'),$$

matrix element have a form:

$$\begin{aligned} M^{\gamma e \to e(n+2)\pi^0} \;&=\; (4\pi\alpha)^{1/2}\left(\frac{\alpha}{\pi f_\pi}\right)^{n+2} sN_e\left[\vec{e}\vec{n}_1\right]\left[\vec{n}_1\vec{n}_2\right]\cdots\left[\vec{n}_{n+1}\vec{n}_{n+2}\right]\times \\ &\times\; R_{01}R_{12}\cdots R_{n+1,n+2}\frac{|\vec{q}_{n+2}|}{q_{n+2}^2}, \end{aligned} \tag{43}$$

the phase volumes of final state is:

$$\begin{aligned} d\Gamma^{\gamma e \to e(n+2)\pi^0} \;&=\; \frac{d^3p_1}{2E_1}\frac{d^3p_2}{2E_2}\cdots\frac{d^3p_{n+2}}{2E_{n+2}}\frac{d^3p'}{2E'}\left(2\pi\right)^{4-3\cdot(n+3)}\times \\ &\times\; \delta^4\left(p+k-p'-p_1-p_2-\cdots-p_{n+2}\right) = \\ &=\; \left(2\pi\right)^{4-3(n+3)}\left(\prod_{i=1}^{n+1}\frac{d\beta_i}{\beta_i}\right)\prod_{i=1}^{n+2}\frac{d^2q_i}{\pi}\frac{1}{2^{n+2}s}\pi^{n+2}, \end{aligned} \tag{44}$$

and the cross section then reads as:

$$\begin{aligned} d\sigma^{\gamma e \to e(n+2)\pi^0} \;&=\; \alpha\left(\frac{\alpha^2}{f_\pi^2}\right)^{n+2}\frac{\pi}{\left(2\pi\right)^{4n+7}}\left(\prod_{i=1}^{n+1}\frac{d^2q_i}{\pi}\right)\frac{\vec{q}_{n+2}^{\,2}\left(d^2q_{n+2}/\pi\right)}{\left[\vec{q}_{n+2}^{\,2}+m_e^2\left(\frac{M_\pi^4}{s^2\beta_{n+1}^2}\right)\right]^2}\times \\ &\times\; \left[\vec{e}_1\vec{n}_1\right]^2\left[\vec{n}_1\vec{n}_2\right]^2\cdots\left[\vec{n}_{n+1}\vec{n}_{n+2}\right]^2 R_{01}^2R_{12}^2\cdots R_{n+1,n+2}^2\left(\prod_{i=1}^{n+1}\frac{d\beta_i}{\beta_i}\right). \end{aligned} \tag{45}$$

In particular in case of one and two pions production we have:

$$d\sigma^{\gamma e \to e\pi^0} \;=\; \frac{\alpha^3}{f_\pi^2}\frac{1}{8\pi^2}\frac{d^2q}{\pi}\frac{\vec{q}_n^{\,2}}{\left[\vec{q}^{\,2}+m_e^2\left(\frac{M_\pi^2}{s}\right)^2\right]^2}\left[\vec{e}\vec{n}\right]^2 R_{01}^2, \tag{46}$$

$$d\sigma^{\gamma e \to e2\pi^0} \;=\; \frac{\alpha^5}{f_\pi^4}\frac{1}{2^7\pi^6}\frac{d^2q_1d^2q_2}{\pi^2}\frac{\vec{q}_2^{\,2}}{\left[\vec{q}_2^{\,2}+m_e^2\left(\frac{M_\pi^2}{s\beta_1}\right)^2\right]^2}\left[\vec{e}\vec{n}_1\right]^2\left[\vec{n}_1\vec{n}_2\right]^2 R_{01}^2R_{12}^2\frac{d\beta_1}{\beta_1}. \tag{47}$$

In particular the single pion production total cross section is (Primakoff process):

$$\sigma^{\gamma e \to e\pi^0}(s) \;=\; \frac{\alpha^3}{4\pi^2 f_\pi^2}\left(1-\left(\vec{e}\vec{n}\right)\left(\vec{e}^{\,*}\vec{n}\right)\right)\ln\frac{s}{m_eM_\pi}. \tag{48}$$

In case of production of n pions on electron-electron collisions:

$$e(k) + e(p) \;\to\; e(k') + e(p') + \pi^0(p_1) + \pi^0(p_2) + \cdots + \pi^0(p_n),$$

the matrix element have a form:

$$M^{ee\to ee(n\cdot\pi^0)} = 2sN_1N_2\left(4\pi\alpha\right)\left(\frac{\alpha}{\pi f_\pi}\right)^n\left(\prod_{i=1}^{n}\left[\vec{n}_i\vec{n}_{i+1}\right]R_{i,i+1}\right) \tag{49}$$

the phase volumes of final state is:

$$\begin{aligned} d\Gamma^{ee\to ee(n\cdot\pi^0)} &= \frac{d^3k'}{2E'_k}\frac{d^3p'}{2E'_p}\frac{d^3p_1}{2E_1}\frac{d^3p_2}{2E_2}\cdots\frac{d^3p_n}{2E_n}\left(2\pi\right)^{4-3\cdot(n+2)}\times \\ &\times\ \delta^4\left(p+k-p'-k'-p_1-p_2-\cdots-p_n\right)= \\ &= \left(2\pi\right)^{4-3(n+2)}\frac{1}{2^{n+1}s}\pi^{n+1}\prod_{i=1}^{n+1}\frac{d^2q_i}{\pi}\prod_{i=1}^{n}\frac{d\beta_i}{\beta_i}, \end{aligned} \tag{50}$$

and the cross section then reads as:

$$\begin{aligned} d\sigma^{ee\to ee(n\cdot\pi^0)} &= \frac{\alpha^{2(1+n)}}{2^{4n-2}\pi^{4n-1}f_\pi^{2n}}\left(\prod_{i=2}^{n}\frac{d^2q_i}{\pi}\right)\left(\prod_{i=1}^{n}\left[\vec{n}_i\vec{n}_{i+1}\right]^2R_{i,i+1}^2\right)\times \\ &\times\ \frac{d^2q_1}{\pi}\frac{\vec{q}_1^{\,2}}{\left[\vec{q}_1^{\,2}+m_1^2\beta_1^2\right]^2}\frac{d^2q_{n+1}}{\pi}\frac{\vec{q}_{n+1}^{\,2}}{\left[\vec{q}_{n+1}^{\,2}+m_2^2\alpha_{n+1}^2\right]^2}\left(\prod_{i=1}^{n}\frac{d\beta_i}{\beta_i}\right), \end{aligned} \tag{51}$$

where $\alpha_{n+1} = M_\pi^2/(s\beta_n)$. In particular in case of one and two pions production we have:

$$d\sigma^{ee\to ee\pi^0} = \frac{\alpha^4}{4\pi^3f_\pi^2}\frac{d^2q_1}{\pi}\frac{\vec{q}_1^{\,2}}{\left[\vec{q}_1^{\,2}+m_1^2\beta_1^2\right]^2}\frac{d^2q_2}{\pi}\frac{\vec{q}_2^{\,2}}{\left[\vec{q}_2^{\,2}+m_2^2\alpha_2^2\right]^2}\left[\vec{n}_1\vec{n}_2\right]^2R_{12}^2\frac{d\beta_1}{\beta_1}, \tag{52}$$

$$\begin{aligned} d\sigma^{ee\to ee2\pi^0} &= \frac{\alpha^6}{2^6\pi^7f_\pi^4}\frac{d^2q_1}{\pi}\frac{\vec{q}_1^{\,2}}{\left[\vec{q}_1^{\,2}+m_1^2\beta_1^2\right]^2}\frac{d^2q_3}{\pi}\frac{\vec{q}_3^{\,2}}{\left[\vec{q}_3^{\,2}+m_2^2\alpha_3^2\right]^2}\times \\ &\times\ \left[\vec{n}_1\vec{n}_2\right]^2\left[\vec{n}_2\vec{n}_3\right]^2\frac{d^2q_2}{\pi}R_{12}^2R_{23}^2\frac{d\beta_1}{\beta_1}\frac{d\beta_2}{\beta_2}. \end{aligned} \tag{53}$$

In general for process $e+e\to e+e+T$ with T is neutral pion or a single pair in case then the transfers momenta to the projectiles do not exceed some values Q_1, Q_2 $\sqrt{-q_1^2}<Q_1$ and $\sqrt{-q_2^2}<Q_2$ the integral in r.h.s. can be performed with logarithmic accuracy:

$$\int\limits_{M_T^2/s}^{1}\frac{d\beta_1}{\beta_1}\ln\left(\frac{Q_1^2}{m_e^2\beta_1^2}\right)\ln\left(\frac{Q_2^2}{m_e^2\alpha_2^2}\right) = \frac{2}{3}[L_T^2+3L_T\ln\frac{Q_1Q_2}{m_e^2}+6\ln\frac{Q_1}{m_e}\ln\frac{Q_2}{m_e}], \tag{54}$$

where $L_T = \ln\left(s/M_T^2\right)$, $s\alpha_2\beta_1 = M_T^2$ and M_T is the characteristic mass of produced system T. In case of processes for process $p+p\to p+p+T$ the $m_e \to M_p$ replacement should be done. In particular the total cross section of single neutral pion production is

$$\sigma^{ee\to ee\pi^0} = \frac{\alpha^4L_\pi}{12\pi^3f_\pi^2}\left[L_\pi^2+6l\left(L_\pi+l\right)\right], \qquad Q_1=Q_2<M_\pi, \tag{55}$$



FIG. 2: Feynman diagram for n pions pairs production at photon-photon collisions.

where $l = \ln(M_\pi/m_e)$.

IV. THE MULTIPLE PAIR PRODUCTION

Let us now consider the processes of multiple pairs production at electron-positron, electron-photon and photon-photon colliders.

A. Process $\gamma\gamma \to n(e^+e^-)$ or $n(\pi^+\pi^-)$

The process of n pairs production at photon-photon collisions are shown on Fig. 2 and the matrix elements of two, three and n pairs production is shown below:

$$\begin{aligned} M_{(2)}^{(\gamma\gamma)} &= \frac{(4\pi\alpha)^2 s}{q_1^2} 2\Phi_0^A \Phi_0^B, \\ M_{(3)}^{(\gamma\gamma)} &= \frac{(4\pi\alpha)^3 s}{q_1^2 q_2^2} 2^2 \Phi_0^A \Phi_{12} \Phi_0^B, \\ &\cdots \\ M_{(2+n)}^{(\gamma\gamma)} &= \frac{(4\pi\alpha)^{n+2} s}{q_1^2 \cdots q_{n+1}^2} 2^{n+1} \Phi_0^A \Phi_{12} \cdots \Phi_{n,n+1} \Phi_0^B, \end{aligned} \tag{56}$$

where Φ_0^A and Φ_0^B are boundary factors (see (19), (20)), which describe pair top and bottom pairs production from one real and one virtual photons. The indexes A and B defines the type of particles which are produced. Factors Φ_{ij} (see (11), (12)) describe pair production in collision of two virtual photons with momenta q_i and q_j. Factor 2^{n+1} comes from Gribov's representation of photon propagator (5).

The phase volumes for two, three, and $(n + 2)$ pairs production have a form (by analogy with the case of two pairs production, considered above, we introduce one, two, and $(n+1)$ transferred momenta):

$$\begin{aligned}
d\Gamma^{(\gamma\gamma)}_{(2)} &= 2^{-9}\pi^{-5}\frac{1}{s}\left(\frac{ds_1 d\Gamma^{\pm}_1}{\pi}\frac{ds_2 d\Gamma^{\pm}_2}{\pi}\right)\frac{d\vec{q}_1}{\pi}, \\
d\Gamma^{(\gamma\gamma)}_{(3)} &= 2^{-16}\pi^{-9}\frac{1}{s}\left(\frac{ds_1 d\Gamma^{\pm}_1}{\pi}\frac{ds_2 d\Gamma^{\pm}_2}{\pi}\frac{ds_3 d\Gamma^{\pm}_3}{\pi}\right)\frac{d\vec{q}_1}{\pi}\frac{d\vec{q}_2}{\pi}\frac{d\beta_1}{\beta_1}, \\
\cdots & \\
d\Gamma^{(\gamma\gamma)}_{(2+n)} &= 2^{-7n-9}\pi^{-4n-5}\frac{1}{s}\left(\prod_{i=1}^{n+2}\frac{ds_i d\Gamma^{\pm}_i}{\pi}\right)\left(\prod_{i=1}^{n+1}\frac{d\vec{q}_i}{\pi}\right)\left(\prod_{i=1}^{n}\frac{d\beta_i}{\beta_i}\right), \qquad (57)
\end{aligned}$$

where s_i and $d\Gamma^{\pm}_i$ are the invariant mass and phase volume of i-th pair (14).

And the cross section for $(2+n)$ pairs production have a form:

$$\begin{aligned}
\sigma^{(\gamma\gamma)}_{(2+n)} &= \frac{1}{8s}\int\left|M^{(\gamma\gamma)}_{(2+n)}\right|^2 d\Gamma^{(\gamma\gamma)}_{(2+n)} = \\
&= \frac{\alpha^{2(n+2)}}{2^{n+2}\pi^{2n+1}}\int\left(\prod_{i=1}^{n+1}\frac{d\vec{q}_i}{\pi\left(\vec{q}_i^{\,2}\right)^2}\right)\left(\prod_{i=1}^{n}\frac{d\beta_i}{\beta_i}\right) I^A_{01}J_{12}J_{23}\cdots J_{n,n+1}I^B_{n+1,0}, \qquad (58)
\end{aligned}$$

where we entered the following impact factors:

$$\begin{aligned}
I^A_{01} &= \int\frac{s_1 d\Gamma^{\pm}_1}{\pi}\left|\Phi^A\right|^2, \qquad J_{ij} = \int\frac{s_i d\Gamma^{\pm}_i}{\pi}\left|\Phi_{ij}\right|^2, \\
I^B_{n+1,0} &= \int\frac{s_{n+2} d\Gamma^{\pm}_{n+2}}{\pi}\left|\Phi^B\right|^2 .
\end{aligned}$$

The explicit form of them was already calculated before (see (15), (16), (17), (21), (22)).

B. Process $\gamma e \to n(e^+e^-)e$ or $n(\pi^+\pi^-)e$

The process of n pairs production at electron-photon collisions are shown on Fig. 3 and the matrix elements of two, three and $2+n$ pairs production is shown below:

$$\begin{aligned}
M^{(\gamma e)}_{(1)} &= \frac{(4\pi\alpha)^{\frac{3}{2}}s}{q_1^2}2N_B\Phi^A_0, \\
M^{(\gamma e)}_{(2)} &= \frac{(4\pi\alpha)^{\frac{5}{2}}s}{q_1^2 q_2^2}2^2N_B\Phi^A_0\Phi_{12}, \\
\cdots & \\
M^{(\gamma e)}_{(2+n)} &= \frac{(4\pi\alpha)^{\frac{2n+5}{2}}s}{q_1^2\cdots q_{n+2}^2}2^{n+2}N_B\Phi^A_0\Phi_{12}\cdots\Phi_{n+1,n+2}, \qquad (59)
\end{aligned}$$

where N_B is the boundary factor, which describe bottom electron line. The phase volumes for one, two, and $(n+2)$ pairs production have a form: (again we introduce



FIG. 3: Feynman diagram for $n+2$ pairs production at electron-photon collisions.

the set of transferred momenta)

$$\begin{aligned}
d\Gamma^{(\gamma e)}_{(1)} &= 2^{-6}\pi^{-3}\frac{1}{s}\left(\frac{ds_1 d\Gamma_1^{\pm}}{\pi}\right)\frac{d\vec{q}_1}{\pi}, \\
d\Gamma^{(\gamma e)}_{(2)} &= 2^{-13}\pi^{-7}\frac{1}{s}\left(\frac{ds_1 d\Gamma_1^{\pm}}{\pi}\frac{ds_2 d\Gamma_2^{\pm}}{\pi}\right)\frac{d\vec{q}_1}{\pi}\frac{d\vec{q}_2}{\pi}\frac{d\beta_1}{\beta_1}, \\
\cdots \\
d\Gamma^{(\gamma e)}_{(2+n)} &= 2^{-7n-13}\pi^{-4n-7}\frac{1}{s}\left(\prod_{i=1}^{n+2}\frac{ds_i d\Gamma_i^{\pm}}{\pi}\right)\left(\prod_{i=1}^{n+2}\frac{d\vec{q}_i}{\pi}\right)\left(\prod_{i=1}^{n+1}\frac{d\beta_i}{\beta_i}\right),
\end{aligned} \tag{60}$$

and the cross section for $(2+n)$ pairs production have a form:

$$\begin{aligned}
\sigma^{(\gamma e)}_{(2+n)} &= \frac{1}{8s}\int\sum\left|M^{(\gamma e)}_{(2+n)}\right|^2 d\Gamma^{(\gamma e)}_{(2+n)} = \\
&= \frac{\alpha^{2n+5)}}{2^n\pi^{2(n+1)}}\int\left(\prod_{i=1}^{n+1}\frac{d\vec{q}_i}{\pi\left(\vec{q}_i^{\,2}\right)^2}\right)\left(\prod_{i=1}^{n+1}\frac{d\beta_i}{\beta_i}\right)\left(2\ln\left(\frac{s\beta_{n+1}}{m^2}\right)\right)\times \\
&\times\ I^A_{01}J_{12}\cdots J_{n,n+1}I_{n+1,0}, \qquad I_{n+1,0} = \left(\frac{J_{n+1,n+2}}{\vec{q}_{n+2}^{\,2}}\right)_{\vec{q}_{n+2}=0}.
\end{aligned} \tag{61}$$

For processes $\gamma e \to eT$ the Weizsaecker-Williams enhancement mechanism works. As a result the factors $\ln(1/\alpha^2_{n+2}) \approx \ln(s\beta_{n+1}/M_T^2)^2$, $\ln(1/\beta^2)$ appears. For details see Appendix A.

In particular cases of creation of one and two fermion pairs we obtain (in agreement with Lipatov and Frolov result [1, 2, 3]):

$$\sigma^{\gamma e\to e^+e^-e} = \frac{28}{9}\frac{\alpha^3}{m^2}L_e,$$



FIG. 4: Feynman diagram for $n+2$ pairs production at electron-positron collisions.

$$\sigma^{\gamma e\to 2e^+2e^-e} = \frac{\alpha^5}{\pi^2 m^2} L_e^2 J_e, \qquad Q_2 < m, \tag{62}$$

where $L_e = \ln\left(s/m_e^2\right)$ and J_e was defined in (26). In cases of one and two charged pion pairs production we obtain:

$$\begin{aligned}\sigma^{\gamma e\to \pi^+\pi^- e} &= \frac{2\alpha^3}{9M_\pi^2} \ln\left(\frac{s}{mM_\pi}\right), \qquad Q_2 < M_\pi, \\ \sigma^{\gamma e\to 2\pi^+2\pi^- e} &= \frac{\alpha^5}{\pi^2 M_\pi^2} L_\pi L_e J_\pi, \qquad Q_2 < M_\pi,\end{aligned} \tag{63}$$

where $L_\pi = \ln\left(s/M_\pi^2\right)$ and J_π was defined in (28).

C. Process $e^+e^- \to n(e^+e^-)$ or $n(\pi^+\pi^-)$

The process of n pairs production at electron-positron collisions are shown on Fig. 4. The matrix elements of one and $(2+n)$ pairs production is shown below:

$$\begin{aligned}M_{(1)}^{(ee)} &= \frac{(4\pi\alpha)^2 s}{q_1^2 q_2^2} 2^2 N_A \Phi_{12} N_B, \\ &\dots \\ M_{(2+n)}^{(ee)} &= \frac{(4\pi\alpha)^{n+3} s}{q_1^2 \cdots q_{n+3}^2} 2^{n+3} N_A \Phi_{12} \cdots \Phi_{n+2,n+3} N_B,\end{aligned} \tag{64}$$

where N_A, N_B are lepton's impact factors. The phase volumes for one, two and $(n+2)$ pairs production have a form:

$$\begin{aligned}
d\Gamma^{(ee)}_{(1)} &= 2^{-10}\pi^{-5}\frac{1}{s}\left(\frac{ds_1 d\Gamma_1^{\pm}}{\pi}\right)\frac{d\vec{q}_1}{\pi}\frac{d\vec{q}_2}{\pi}\frac{d\beta_1}{\beta_1}, \\
d\Gamma^{(ee)}_{(2)} &= 2^{-17}\pi^{-9}\frac{1}{s}\left(\frac{ds_1 d\Gamma_1^{\pm}}{\pi}\frac{ds_2 d\Gamma_2^{\pm}}{\pi}\right)\frac{d\vec{q}_1}{\pi}\frac{d\vec{q}_2}{\pi}\frac{d\vec{q}_3}{\pi}\frac{d\beta_1}{\beta_1}\frac{d\beta_2}{\beta_2}, \\
\dots \\
d\Gamma^{(ee)}_{(2+n)} &= 2^{-7n-17}\pi^{-4n-9}\frac{1}{s}\left(\prod_{i=1}^{n+2}\frac{ds_i d\Gamma_i^{\pm}}{\pi}\right)\left(\prod_{i=1}^{n+3}\frac{d\vec{q}_i}{\pi}\right)\left(\prod_{i=1}^{n+2}\frac{d\beta_i}{\beta_i}\right),
\end{aligned} \tag{65}$$

and the cross section for $(2+n)$ pairs production have a form

$$\begin{aligned}
\sigma^{(ee)}_{(2+n)} &= \frac{1}{8s}\int\sum\left|M^{(ee)}_{(2+n)}\right|^2 d\Gamma^{(ee)}_{(2+n)} = \\
&= \frac{\alpha^{2(n+3)}}{2^{n-2}\pi^{2n+3}}\int\left(\prod_{i=1}^{n+1}\frac{d\vec{q}_i}{\pi\left(\vec{q}_i^{\,2}\right)^2}\right)\left(\prod_{i=1}^{n+2}\frac{d\beta_i}{\beta_i}\right)\left(2\ln\left(\frac{1}{\beta_1}\right)\right)\left(2\ln\left(\frac{s\beta_{n+2}}{m^2}\right)\right)\times \\
&\times I_{01}\left(\prod_{i=1}^{n+1}J_{i,i+1}\right)I_{n+2,0}.
\end{aligned} \tag{66}$$

For the case of creation one and two electron-positron pairs we reproduce the known results (see [1, 2, 3, 4, 5, 6, 9, 10]):

$$\begin{aligned}
\sigma^{ee\to eee^+e^-} &= \frac{28}{27}\frac{\alpha^4 L_e^3}{\pi m^2}, \\
\sigma^{ee\to ee2e^+2e^-} &= \frac{\alpha^6 L_e^4}{6\pi^3 m^2}J_e, \qquad Q_1\sim Q_2<m.
\end{aligned} \tag{67}$$

For processes $e\bar{e}\to e\bar{e}\pi^+\pi^-$ and $e\bar{e}\to e\bar{e}2\pi^+2\pi^-$ we obtain:

$$\begin{aligned}
\sigma^{e\bar{e}\to e\bar{e}\pi^+\pi^-} &= \frac{4}{27}\frac{\alpha^4 L_\pi}{M_\pi^2}[L_\pi^2+6l(L_\pi+l)], \qquad Q_1\sim Q_2<m, \\
\sigma^{e\bar{e}\to e\bar{e}2\pi^+2\pi^-} &= \frac{\alpha^6 L_\pi^2}{6\pi^3 M_\pi^2}\left(L_\pi^2+8lL_\pi+12l^2\right)J_\pi, \qquad Q_1\sim Q_2<m,
\end{aligned}$$

where $L_\pi=\ln\left(s/M_\pi^2\right)$ and $l=\ln\left(M_\pi/m_e\right)$.

V. GENERALIZATION

Let us pay some attention to obtained cross sections (58), (61), (67). They have similar form. The integration over β_i leads to some logarithmical enhancement. Let us demonstrate this in case of electron-positron pairs production. Keeping in mind the relative arrangements of β_i (see (35)) we obtain:

$$\int \prod_{i=1}^{n} \frac{d\beta_i}{\beta_i} = L_e^n \int_0^1 dx_1 \int_0^{x_1} dx_2 \cdots \int_0^{x_{n-1}} dx_n = \frac{L_e^n}{n!}. \tag{68}$$

This can be presented as the integral over complex variable j:

$$\frac{L^n}{n!} = \oint_C \frac{dj}{2\pi i} \frac{\left(s/m^2\right)^j}{j^{n+1}}, \tag{69}$$

where contour C is the small circle around the point $j = 0$ in the j complex plane. Denoting the common part as:

$$Z_n = \frac{\alpha^{2n}}{2^n \pi^{2n} j^n} \left(\prod_{i=1}^{n} J_{i,i+1} \right) \left(\int \prod_{i=1}^{n+1} \frac{d\vec{q}_i}{\pi \left(\vec{q}_i^{\,2}\right)^2} \right), \tag{70}$$

we can rewrite these cross sections in form:

$$\begin{aligned} \sigma_{(2+n)}^{(\gamma\gamma)} &= \oint \frac{dj}{2\pi i} \left(s/m^2\right)^j \left(\frac{\alpha^4}{4\pi} I_{01}^A Z_n I_{n+1,0}^B \right), \\ \sigma_{(2+n)}^{(\gamma e)} &= \oint \frac{dj}{2\pi i} \left(s/m^2\right)^j \left(\frac{\alpha^5}{\pi^2 j^2} I_{01}^A Z_n I_{n+1,0} \right), \\ \sigma_{(2+n)}^{(ee)} &= \oint \frac{dj}{2\pi i} \left(s/m^2\right)^j \left(\frac{4\alpha^6}{\pi^3 j^4} I_{01} Z_n I_{n+1,0} \right). \end{aligned} \tag{71}$$

Considering the production of 1, 2, ... pairs it is convenient to introduce the value Ψ:

$$\Psi = \sum_{n=1}^{\infty} Z_n = \sum_{n=1}^{\infty} \frac{\alpha^{2n}}{2^n \pi^{2n} j^n} \left(\prod_{i=1}^{n} J_{i,i+1} \right) \left(\int \prod_{i=1}^{n+1} \frac{d\vec{q}_i}{\pi \left(\vec{q}_i^{\,2}\right)^2} \right). \tag{72}$$

Then the cross sections of production of 1, 2, ... pairs of charged particles reads as:

$$\begin{aligned} \sigma_{n\geq 3}^{(\gamma\gamma)} &= \oint \frac{dj}{2\pi i} \left(s/m^2\right)^j \left(\frac{\alpha^4}{4\pi} I_{01}^A \Psi I_{n+1,0}^B \right), \\ \sigma_{n\geq 1}^{(\gamma e)} &= \oint \frac{dj}{2\pi i} \left(s/m^2\right)^j \left(\frac{\alpha^5}{\pi^2 j^2} I_{01}^A \Psi I_{n+1,0} \right), \\ \sigma_{n\geq 1}^{(ee)} &= \oint \frac{dj}{2\pi i} \left(s/m^2\right)^j \left(\frac{4\alpha^6}{\pi^3 j^4} I_{01} \Psi I_{n+1,0} \right). \end{aligned} \tag{73}$$

VI. KINEMATICS OF JETS EMISSION

Let discuss the kinematics of the definite pair production at collision of two virtual photons:

$$\gamma^*(q_i) + \gamma^*(-q_{i+1}) \rightarrow e^+(q_{+i}) + e^-(q_{-i}), i = 1, 2, ..., n.$$

Keeping in mind the Sudakov parametrization of the 4-momentum of pairs component $q^i_\pm = \alpha_{\pm i} E n_p + \beta_{\pm i} E n_k + q_{\pm\perp i}$ with $2E = \sqrt{s}$ is the center of mass beams total energy. Light-like 4 vectors $n_p = (1,1,0,0)$, $n_k = (1,-1,0,0)$ have space component-unit vectors directed along initial 3-momenta of colliding beams, $\vec{p}$, $\vec{k}$. We choice a vector $\vec{k}$ as z-axes direction. The transverse to the beam direction component $q_\perp = (0,0,q_x,q_y)$, $q^2_\perp = -\vec{q}^2$ can be expressed in terms of the polar angle of the relevant particle emission in the center of mass frame:

$$\tan\theta_i = \frac{|\vec{q}_i|}{E(\beta_i - \alpha_i)}, \qquad \alpha_i = \frac{\vec{q}_i^2 + m^2}{4E^2\beta_i}, \qquad i = \pm. \tag{74}$$

Particle with momentum q_i moves in hemisphere $\vec{p}$ if $\alpha_i > \beta_i$ and to hemisphere $\vec{k}$ in case $\beta_i > \alpha_i$. We remind here the conservation low $q_i - q_{i+1} = q_{-i} + q_{+i}$. In Sudakov parametrization it can be written as

$$\vec{q}_{\perp,i} - \vec{q}_{\perp,i+1} = \vec{q}_{+i} + \vec{q}_{-i}, \quad \beta_i = \beta_{+i} + \beta_{-i}, \quad -\alpha_{i+1} = \alpha_{+i} + \alpha_{-i}, \tag{75}$$

where we keep into account the rigorous arrangements of longitudinal Sudakov parameters:

$$\begin{aligned} &1 \gg \beta_1 \gg \beta_2 \gg \cdots \gg \beta_n \gg \frac{m_1^2}{E^2}, \\ &1 \gg \alpha_n \gg \alpha_{n-1} \gg \cdots \gg \alpha_1 \gg \frac{m_2^2}{E^2}, \end{aligned} \tag{76}$$

with $m_{1,2}$ are the masses of colliding particles or the mass of the created pair component in the case when the relevant initial particle is a photon.

In paper [11] this results were generalized to the supersymmetric QED case.

VII. INCLUSIVE CROSS SECTION

In the experimental set-up with detecting only one pair emitted in peripheral kinematics the differential distributions averaged on kinematical parameters of other pairs. Let us describe the modification of cross section of production of n pairs to obtain the contribution to inclusive cross section. Keeping in mind the rigorous arrangement by magnitude of longitudinal Sudakov parameters we can conclude that two replacements must be done. One of them concern the distribution on the longitudinal Sudakov parameters of the momentum transferred. Really the replacement must be done: (m is the mass of pair components)

$$\frac{\delta}{\delta\beta_k} \int\limits_{m^2/s}^{1} \frac{d\beta_1}{\beta_1} \int\limits_{m^2/s}^{\beta_1} \frac{d\beta_2}{\beta_2} \cdots \int\limits_{m^2/s}^{\beta_{n-1}} \frac{d\beta_n}{\beta_n} =$$

$$= \frac{1}{\beta_k}\ln^{k-1}(1/\beta_k)(L_e - \ln(1/\beta_k))^{n-k}\frac{1}{\Gamma(k)\Gamma(n-k+1)} \equiv F_{k,n}(\beta_k). \quad (77)$$

The quantity $F_{k,n}(\beta)$ is normalized as

$$\int\limits_{m^2/s}^{1} d\beta F_{k,n}(\beta) = \frac{L_e^n}{n!}. \quad (78)$$

Another one is

$$G_{k,n}(\vec{\Delta}^2) \equiv \pi\frac{\delta^2}{\delta^2\Delta}\int\left(\prod_{i=1}^{n+1}\frac{d\vec{q}_i}{\pi\left(\vec{q}_i^2\right)^2}\right)\left(\prod_{i=1}^{n}\frac{d\beta_i}{\beta_i}\right)I_{01}^{A}J_{12}J_{23}\cdots J_{n,n+1}I_{n+1,0}^{B}, \quad (79)$$

where $\vec{\Delta} = \vec{q}_k - \vec{q}_{k+1}$. Keeping these factors together we obtain:

$$m^4\pi\frac{d\sigma^{\gamma\gamma}}{d\beta d^2\Delta} = \sum_{1}^{\infty}\frac{\alpha^{2(n+2)}}{4\pi(2\pi^2)^n}F_{k,n}(\beta)G_{k,n}(x), \qquad x = \frac{\vec{\Delta}^2}{m^2}. \quad (80)$$

The lowest order inclusive cross section of lepton pair production in $\gamma\gamma$ collisions for the case of unpolarized photon beams is:

$$m^4\beta\pi\frac{d\sigma_1^{\gamma\gamma}}{d\beta d^2\Delta} = \left(\frac{\alpha^2}{2\pi}\right)^3 G_1(x)L_e,$$

$$\begin{aligned} G_1(x) &= \frac{16}{9}m^4\int\frac{d^2q}{\pi}\int\limits_0^1 dx_1\int\limits_0^1 dx_2\int\limits_0^1 dx_3\int\limits_0^1 dx_4 \times \\ &\times\left(\frac{1+a_1}{m^2+a_1\vec{q}^2}\right)\left(\frac{1+a_2}{m^2+a_2(\vec{q}+\vec{\Delta})^2}\right)\left(\frac{a_3+a_4-5a_3a_4}{m^2+a_3\vec{q}^2+a_4(\vec{q}+\vec{\Delta})^2}\right), \end{aligned} \quad (81)$$

$$\begin{aligned} a_1 &= x_1(1-x_1), \qquad a_3 = x_3(1-x_3), \\ a_2 &= x_2(1-x_2), \qquad a_4 = x_4(1-x_4), \end{aligned}$$

where $x = \Delta^2/m^2$. Numerical value of the dimensionless quantity $G_1(x)$ is given in Fig. 5. Numerical integration of this function over all possible tranverse momentum gives

$$\int\limits_0^\infty dx G_1(x) \approx 26.5. \quad (82)$$



FIG. 5: Function $G_1(x)$ (81) dependence of x.

VIII. SUMMATION ON THE PAIR NUMBER

Using representation of logarithmic enhancement factor $L_e^n/n!$ in the form of the integral over complex variable (see (69)) the cross section of arbitrary number of pair production in γe, $\gamma\gamma$ and ee collisions may be written down in the universal form (73), where function $\Psi_j(x_1, x_2)$ is defined as:

$$\Psi_j(x_1, x_2) = \sum_{n=1}^{\infty} h^{-n} \left(\prod_{i=1}^{n} \int \frac{d^2 q_i}{\pi \left(q_i^2\right)^2} \right) J_{12} \cdots J_{n-1,n}. \tag{83}$$

$$h^{-1} = \frac{\alpha^2}{2\pi^2 j}. \tag{84}$$

This function are based on the integral equation:

$$\Psi_j(x_1, x_2) = k(x_1, x_2) + h^{-1} \int_0^{\infty} dx_1' k(x_1, x_1') \Psi_j(x_1', x_2). \tag{85}$$

There are two different contribution to kernel $k(x_1,x_2)$ (we do not consider here π^0 channels): $k(x_1,x_2)=k_e(x_1,x_2)+k_\pi(x_1,x_2)$, with

$$k_e(x_1,x_2) = 4\int\limits_0^1 d\alpha\int\limits_0^1 d\beta\frac{a+b-5ab}{ax_1+bx_2+m^2}, \tag{86}$$

$$k_\pi(x_1,x_2) = 2\int\limits_0^1 d\alpha\int\limits_0^1 d\beta\frac{1/4-(a+b-5ab)}{ax_1+bx_2+m^2}. \tag{87}$$

The function $\Psi_j(x_1,x_2)$ dependence of variable j have a square root singularity in j-plane. To find the most right position of this cut in j-plane we may investigate the homogenous integral equation:

$$\Delta\Psi_j^i(x_1,x_2)=\frac{1}{\kappa^i}\int\limits_0^\infty dx_1' k^i(x_1,x_1')\Delta\Psi_j^i(x_1',x_2), \qquad i=e,\pi. \tag{88}$$

Further simplification consists in putting the mass of created particles equal to zero $m=0$.

To solve this homogeneous integral equation let expand function $\Psi_j(x_1,x_2)$ by full set of functions $(x_1/x_2)^\lambda/\sqrt{x_1x_2}$, with λ pure imaginary parameter and find the relevant eigenvalue..

First we can perform the integration on the intermediate photon momentum. We have:

$$\int\limits_0^\infty\frac{t^\gamma dt}{a+bt}=\frac{\pi}{\sin(\pi\gamma)}a^\gamma b^{-\gamma-1}, \qquad a,b>0, \tag{89}$$

with $\gamma=-1/2+i\eta$. Using the explicit form of kernels k_e, k_π (see (86), (87)) and performing α, β integration after some algebra we obtain:

$$\kappa^e=\frac{\pi^2(11+12\eta^2)\sinh(\pi\eta)}{16\eta(1+\eta^2)\cosh^2(\pi\eta)};$$
$$\kappa^\pi=\frac{\pi^2(5+4\eta^2)\sinh(\pi\eta)}{32\eta(1+\eta^2)\cosh^2(\pi\eta)}. \tag{90}$$

As a result we obtain the following solution:

$$\Psi_j(x_1,x_2)|_{m^2=0}=\frac{1}{2\pi i}\int\limits_{-i\infty}^{i\infty}d\lambda\left(\frac{x_1}{x_2}\right)^\lambda\frac{1}{\sqrt{x_1x_2}}\left(\frac{1}{\kappa(\lambda)}-\frac{1}{h(j)}\right)^{-1}, \tag{91}$$

where

$$\kappa(\lambda)=\frac{\pi^2\sin(\pi\lambda)}{128\lambda(1-\lambda^2)\cos^2(\pi\lambda)}\left[2N_f\left(11-12\lambda^2\right)+N_b\left(5-4\lambda^2\right)\right], \tag{92}$$

where N_f is the number of charged leptons and N_b is the number of charged mesons. The solution if the analytical function of j with the cut located at $1 < j < j_0 = 1 + \frac{\pi\alpha^2}{64}\left[22N_f + 5N_b\right]$. This formula is the generalization of formula obtained in [1, 2, 3] where only charged fermions case was considered.

IX. CONCLUSION

The partially integrated cross sections, which takes into account the transfer momentum not exceeding $\left|\vec{q}_1^{\,2}\right| < Q_1 = \sqrt{-q_1^2}$, $\left|\vec{q}_2^{\,2}\right| < Q_2 = \sqrt{-q_2^2}$ are given in Appendix B. We choose $L_\pi \approx 20$ which is expected at LHC. It can be seen that the cross sections are big enough to be taken into account and measured.

We should note that the deviation from this asymptotic formulae can be considerable. For instance the cross section of process $pp \to ppe\bar{e}$ (see formula (B4) in Appendix B) gives $\sigma^{LL}_{pp\to ppe\bar{e}} = 9.9$ mb at $\sqrt{s} = 14$ TeV, while the more precise form is [10]:

$$\sigma^{pp\to ppe\bar{e}} = \frac{28}{27}\frac{\alpha^4}{\pi m_e^2}\left(L_p^3 + AL_p^2 + BL_p + C\right), \tag{93}$$

where $L_p = \ln\left(s/M_p^2\right)$ and

$$\begin{aligned} A &= -\frac{178}{28} = -6.35714, \\ B &= \frac{1}{28}\left(7\pi^2 + 370\right) = 15.6817, \\ C &= -\frac{1}{28}\left(348 + \frac{13}{2}\pi^2 - 21\zeta_3\right) = -13.8182, \end{aligned}$$

which gives $\sigma_{pp\to ppe\bar{e}} = 7$ mb. Again for process $e\bar{e} \to 2e2\bar{e}$ we have following precise result [12]:

$$\sigma^{e\bar{e}\to 2e2\bar{e}} = \frac{\alpha^4}{\pi m_e^2}\left(AL_e^3 + BL_e^2 + CL_e + D\right), \tag{94}$$

where $L_e = \ln\left(s/m_e^2\right)$ and

$$\begin{aligned} A &= \frac{28}{27} = 1.03704, \\ B &= -\frac{178}{27} = -6.59259, \\ C &= -\frac{164}{9}\zeta_2 + \frac{490}{27} = -11.8262, \\ D &= \frac{52}{3}\zeta_2 \ln 2 + \frac{401}{9}\zeta_3 + \frac{886}{27}\zeta_2 - \frac{428}{27} = 111.448. \end{aligned}$$

In leading logarithmical approximation it gives $\sigma^{LL}_{e\bar{e}\to 2e2\bar{e}} = 56$ mb, while the precise value is $\sigma_{e\bar{e}\to 2e2\bar{e}} = 45$ mb at $\sqrt{s} = 14$ TeV.

So the formulae given above have the accuracy of order $\pm(15-20)\%$.

X. ACKNOWLEDGEMENTS

Two of us (E.A.K. and Yu.M.B.) are grateful to INTAS grant (05-1000008-8328) and Belorussian grant MK-2952.2006.2 for financial support. We are also grateful to S. Dubnicka, C. Adamuscin, A. Z. Dubnickova for useful discussions. We also grateful to Lev Lipatov for critical comments.

APPENDIX A: WEIZSAECKER-WILLIAMS (WW) ENHANCEMENT FOR γe AND e^+e^- COLLISIONS

Let us consider the details of $e \to \gamma^* e$ vertex

$$e(p_1) \to \gamma^*(q) + e(p_1')$$

in peripherical kinematics. Using Sudakov parametrization of the 4-momentum of virtual photon:

$$q = \alpha p + \beta \tilde{p}_1 + q_\perp, \quad p^2 = \tilde{p}_1^2 = 0, \quad q_\perp p = q_\perp \tilde{p}_1 = 0, \quad 2p_1\tilde{p}_1 = m^2, \tag{A1}$$

and keeping in mind the electrons on mass shell conditions:

$$(p_1 - q)^2 - m^2 = -s\alpha(1-\beta) - m^2\beta - \vec{q}^2 = 0, \tag{A2}$$

we obtain for the square of virtual photon momentum:

$$q^2 = -\frac{1}{1-\beta}\left[\vec{q}^2 + m^2\beta^2\right]. \tag{A3}$$

We use here the standard definitions $q_\perp^2 = -\vec{q}^2$, $s = 2pp_1 \gg m^2, \vec{q}^2$. It is important to note that transfer momentum is space-like vector with nonzero norm. Performing integration on $d^2\vec{q}$ in matrix element square and keeping in mind the consequences of gauge invariance we obtain an enhancement factor in rapidity β:

$$\int\limits_0^{Q^2} \frac{\vec{q}^2 d\vec{q}^2}{(\vec{q}^2 + m^2\beta^2)^2} \approx \ln\frac{Q^2}{m^2\beta^2}. \tag{A4}$$

APPENDIX B: THE TOTAL CROSS SECTIONS OF π_0 AND $\pi_+\pi_-$ PRODUCTION

Let us consider the total cross sections of processes of pions production within logarithmic accuracy. In case then the momentum transfer does not exceed the

$Q_1, Q_2 < M_\pi$ we have (see comments to formula (55)):

$$\begin{aligned}
\sigma^{ee\to ee\pi^0} &= \frac{\alpha^4 L_\pi}{12\pi^3 f_\pi^2}\left[L_\pi^2 + 6\ln\left(\frac{M_\pi}{m_e}\right)\ln\left(\frac{s}{M_\pi m_e}\right)\right], \qquad Q_1 \sim Q_2 < M_\pi,\\
\sigma^{ee\to ee\pi^+\pi^-} &= \frac{4}{27}\frac{\alpha^4}{M_\pi^2}L_\pi\left[L_\pi^2 + 6\ln\left(\frac{M_\pi}{m_e}\right)\ln\left(\frac{s}{M_\pi m_e}\right)\right], \qquad Q_1 \sim Q_2 < M_\pi,\\
\sigma^{pp\to pp\pi^0} &= \frac{\alpha^4 L_\pi^3}{12\pi^3 f_\pi^2}, \qquad Q_1 \sim Q_2 < M_P,\\
\sigma^{ep\to ep\pi^0} &= \frac{\alpha^4 L_\pi^2}{12\pi^3 f_\pi^2}\left[L_\pi + 3\ln\left(\frac{M_\pi}{m_e}\right)\right], \qquad Q_1 < M_\pi, \quad Q_2 < M_P,\\
\sigma^{\gamma p\to p2\pi^0} &= \frac{\alpha^5}{2^8\pi^6 f_\pi^2}\left(L_\pi^2 + L_\pi\ln\left(\frac{M_\pi}{m_e}\right)\right)\int\frac{d^2\vec{q}_1}{\pi f_\pi^2}R_{01}^4, \qquad Q_2 < M_\pi,\\
\sigma^{\gamma e\to e\pi^0} &= \frac{\alpha^3}{4f_\pi^2}\ln\left(\frac{s}{M_\pi m_e}\right), \qquad Q_2 < M_\pi,\\
\sigma^{\gamma e\to e\pi^+\pi^-} &= \frac{2\alpha^3}{9M_\pi^2}\ln\left(\frac{s}{M_\pi m_e}\right), \qquad Q_2 < M_\pi,\\
\sigma^{\gamma e\to e2\pi^0} &= \frac{\alpha^5}{2^8\pi^6 f_\pi^2}\left(L_\pi^2 + 2L_\pi\ln\left(\frac{M_\pi}{m_e}\right)\right)\int\frac{d^2 q_1}{\pi f_\pi^2}R_{01}^4, \qquad Q_2 < M_\pi,\\
\sigma^{\gamma e\to e2(\pi^+\pi^-)} &= \frac{\alpha^5}{\pi^2 M_\pi^2}L_\pi\ln\left(\frac{s}{M_\pi m_e}\right)J_\pi,\\
\sigma^{\gamma\gamma\to 2\pi^0} &= \frac{\alpha^4}{2^8\pi^5 f_\pi^2}\int\frac{d^2\vec{q}_1}{\pi f_\pi^2}R_{01}^4,\\
\sigma^{\gamma\gamma\to 2(\pi^+\pi^-)} &= \frac{\alpha^4}{\pi M_\pi^2}J_\pi,\\
\sigma^{\gamma\gamma\to 3\pi^0} &= \frac{\alpha^6}{2^{13}\pi^9 f_\pi^2}L_\pi\int\frac{d\vec{q}_1^{\,2}}{f_\pi^2}\int\frac{d\vec{q}_2^{\,2}}{f_\pi^2}R_{01}^2 R_{02}^2 R_{12}^2.
\end{aligned}$$

The integrals R_{01}, R_{02}, R_{12} have a form:

$$\begin{aligned}
R_{01}\left(\vec{q}_1^{\,2}\right) &= 2M_q^2\int_0^1 dx_1\int_0^{1-x_1}\frac{dx_2}{M_q^2 - M_\pi^2 x_2 x_3 + \vec{q}_1^{\,2}x_1 x_3}, \qquad x_3 = 1 - x_1 - x_2,\\
R_{12}\left(\vec{q}_1^{\,2}, \vec{q}_2^{\,2}\right) &= 2M_q^2\int_0^1 dx_1\int_0^{1-x_1}\frac{dx_2}{M_q^2 - M_\pi^2 x_2 x_3 + \left(\vec{q}_1^{\,2}x_2 + \vec{q}_2^{\,2}x_3\right)x_1},
\end{aligned} \tag{B1}$$

where M_q is the loop quark mass ($M_q \sim 260$ MeV). We keep into account only the logarithm of ratio of pion and electron masses $\ln\left(M_\pi/m_e\right)$. As rather good approximation (within 5% accuracy) we can omit term $\left(-M_\pi^2 x_2 x_3\right)$ compared with M_q^2 in

R_{01} and R_{12} denominators and obtain the following formulae:

$$R_{01}\left(\vec{q}_1^{\,2}\right) = \frac{M_q^2}{\vec{q}_1^{\,2}} \ln^2\left(\frac{b+1}{b-1}\right), \qquad b = \sqrt{1+\frac{4M_q^2}{\vec{q}_1^{\,2}}}, \tag{B2}$$

$$R_{12}\left(\vec{q}_1^{\,2}, \vec{q}_2^{\,2}\right) = \frac{\vec{q}_1^{\,2} R_{01}\left(\vec{q}_1^{\,2}\right) - \vec{q}_2^{\,2} R_{01}\left(\vec{q}_2^{\,2}\right)}{\vec{q}_1^{\,2} - \vec{q}_2^{\,2}}. \tag{B3}$$

The numerical evaluation of phase integrals gives:

$$\int \frac{d^2\vec{q}_1}{\pi f_\pi^2} R_{01}^4 \approx 46.2$$

$$\int \frac{d^2\vec{q}_1}{\pi f_\pi^2} \int \frac{d^2\vec{q}_2}{\pi f_\pi^2} R_{01}^2 R_{02}^2 R_{12}^2 \approx 2521.$$

The total cross sections in leading logarithmical approximation of processes $pp \to ppe\bar{e}$ and $pp \to pp\pi^+\pi^-$ have a form:

$$\sigma^{pp\to ppe\bar{e}} = \frac{28}{27}\frac{\alpha^4}{\pi m_e^2} \ln^3\left(\frac{s}{M_p^2}\right), \tag{B4}$$

$$\sigma^{pp\to pp\pi^+\pi^-} = \frac{4}{27}\frac{\alpha^4}{\pi M_\pi^2} \ln^3\left(\frac{s}{M_p^2}\right).$$

For completeness we evaluate this cross sections at energy $\sqrt{s} = 14$ TeV (available at Tevatron). In that case $L_e = 34$, $L_\pi = 23$ and the cross sections are:

$\sigma^{ee\to ee\pi^0} = 12$ nb,
$\sigma^{ee\to ee\pi^+\pi^-} = 309$ nb,
$\sigma^{pp\to pp\pi^0} = 4.28$ nb,
$\sigma^{ep\to ep\pi^0} = 7.38$ nb,
$\sigma^{\gamma p\to p2\pi^0} = 0.117$ pb,
$\sigma^{\gamma e\to e\pi^0} = 127$ nb,
$\sigma^{\gamma e\to e\pi^+\pi^-} = 53$ nb,
$\sigma^{\gamma e\to e2\pi^0} = 0.14$ pb,
$\sigma^{\gamma e\to e2(\pi^+\pi^-)} = 3.79$ pb,
$\sigma^{\gamma\gamma\to 2\pi^0} = 0.076$ pb.
$\sigma^{\gamma\gamma\to 2(\pi^+\pi^-)} = 2.47$ pb,
$\sigma^{\gamma\gamma\to 3\pi^0} = 1.64$ ab,
$\sigma^{pp\to ppe\bar{e}} = 9.9$ mb,
$\sigma^{pp\to pp\pi^+\pi^-} = 20$ nb,

APPENDIX C: DETAILS OF J_{12}^e CALCULATION

We put here some details of obtaining J_{12}^e from formula (15). After performing traces it can be written down in the form:

$$\begin{aligned} J_{12}^e = \int_0^1 dx_+ \int \frac{d^2q_+}{\pi} \Bigg\{ & \frac{\left(D_1 - D_2\right)^2 \left(\vec{q}_+^{\,2} + m^2\right)\left(\vec{q}_-^{\,2} + m^2\right)}{\left(D_1 D_2\right)^2} + \\ & + \frac{\vec{q}_1^{\,2}\left(\vec{q}_+^{\,2} + m^2\right)}{D_1^2} + \frac{\vec{q}_1^{\,2}\left(\vec{q}_-^{\,2} + m^2\right)}{D_2^2} + \\ & + 2\left(\vec{q}_1\vec{q}_-\right)\left(\vec{q}_+^{\,2} + m^2\right)\frac{D_1 - D_2}{D_1^2 D_2} - 2\left(\vec{q}_1\vec{q}_+\right)\left(\vec{q}_-^{\,2} + m^2\right)\frac{D_1 - D_2}{D_1 D_2^2} - \end{aligned}$$

$$- \left. 2\frac{\vec{q}_1^{\,2}\left(m^2-(\vec{q}_+\vec{q}_-)\right)+2\,(\vec{q}_+\vec{q}_1)\,(\vec{q}_-\vec{q}_1)}{D_1D_2}\right\} . \tag{C1}$$

Expression in figure brackets explicitly vanishes if $\vec{q}_1 \to 0$ or $\vec{q}_2 \to 0$. Each separated term contain ultraviolet divergence and requires regularization. That is the reason to enter the ultraviolet cut off parameter Λ. For example the contribution of second and third terms are:

$$\int \frac{d^2q_+}{\pi}\left(\frac{\vec{q}_1^{\,2}\left(\vec{q}_+^{\,2}+m^2\right)}{D_1^2}+\frac{\vec{q}_1^{\,2}\left(\vec{q}_-^{\,2}+m^2\right)}{D_2^2}\right) =$$

$$= \vec{q}_1^{\,2}\left[2\ln\frac{\Lambda^2}{D_0}+\frac{\vec{q}_2^{\,2}\left(x_+-x_-\right)^2}{D_0}\right], \tag{C2}$$

where $D_0 = x_+x_- - \vec{q}_2^{\,2} + m^2$. In next terms we join the denominators using Feynman trick:

$$\frac{1}{D_1D_2} = \int\limits_0^1 \frac{dt}{\left(tD_1+\bar{t}D_2\right)^2} = \int\limits_0^1 \frac{dt}{\left(\left(\vec{q}_+-\vec{b}\right)^2+D\right)^2}, \tag{C3}$$

where $\vec{b} = \bar{t}\vec{q}_1 + x_+\vec{q}_2$, and $D = m^2 + t\bar{t}\vec{q}_1^{\,2} + x_+x_-\vec{q}_2^{\,2}$. The following integration on $\vec{q}_+$ is straightforward. The dependence of ultraviolet cut off Λ disappears. The result of expression will contain terms with $\ln(D/m^2)$, D^0, D^{-1}, D^{-2}, D^{-3}. Using the integration by parts

$$\int\limits_0^1 dt\frac{\vec{q}_1^{\,2}(1-2t)f(t)}{D^2} = -\left.\frac{f(t)}{D}\right|_0^1 + \int\limits_0^1 \frac{dt}{D}f'(t),$$

we may express terms containing D^{-2}, D^{-3} in terms with D^{-1}. As a result we get:

$$J_{12}^e = \int\limits_0^1 dx_+\int\limits_0^1 dt\left\{2b(1-4b)\left(\frac{1}{D}-\frac{1}{D_0}\right)\left(\vec{q}_2^{\,2}\right)^2 + \right.$$

$$\left. + \frac{1}{D}\left[\vec{q}_1^{\,2}\vec{q}_2^{\,2}(1+8ab-2b)-8ab\left(\vec{q}_1\vec{q}_2\right)^2\right]\right\}, \tag{C4}$$

where $a = t\bar{t}$ and $b = x_+x_-$. The first term in figure brackets can be written in symmetrical form:

$$\int\limits_0^1 dx_+\int\limits_0^1 dt2b(1-4b)\left(\frac{1}{D}-\frac{1}{D_0}\right)\left(\vec{q}_2^{\,2}\right)^2 =$$

$$= 2\vec{q}_2^{\,2}\int\limits_0^1 dx_+\left(x_+x_-(1-2x_+)\right)\frac{d}{dx_+}\ln\frac{D}{D_0} =$$

$$= -\vec{q}_1^{\,2}\vec{q}_2^{\,2}\int\limits_0^1 \frac{dx_+ dt}{D}\,(1-4a)\,(1-6b)\,. \qquad \text{(C5)}$$

Substituting this result to (C4) we obtain (15).

---

[1] G. V. Frolov, V. N. Gribov, and L. N. Lipatov, Phys. Lett. **B31**, 34 (1970).
[2] V. N. Gribov, L. N. Lipatov, and G. V. Frolov, Sov. J. Nucl. Phys. **12**, 543 (1971).
[3] L. N. Lipatov and G. V. Frolov, Yad. Fiz. **13**, 588 (1971).
[4] H. Cheng and T. T. Wu, Phys. Rev. **182**, 1852 (1969).
[5] H. Cheng and T. T. Wu, Phys. Rev. Lett. **22**, 666 (1969).
[6] H. Cheng and T. T. Wu, Phys. Rev. **D1**, 3414 (1970).
[7] E. A. Kuraev, A. Schiller, and V. G. Serbo, Phys. Lett. **B134**, 455 (1984).
[8] E. A. Kuraev, A. Schiller, and V. G. Serbo, Nucl. Phys. **B256**, 189 (1985).
[9] L. D. Landau and E. M. Lifschitz, Phys. Z. Sowjetunion **6**, 244 (1934).
[10] G. Racah, Nuovo Cim. **13**, 69 (1936).
[11] A. V. Kotikov and L. N. Lipatov, Nucl. Phys. **B582**, 19 (2000), hep-ph/0004008.
[12] E. A. Kuraev and L. N. Lipatov, Yad. Fiz. **16**, 1060 (1972).